



LOOMIS BASIN
Est. 1975
 VETERINARY CLINIC

Canine Vaccination Schedule

Recommended (Core) Vaccines

Disease	Description	Primary puppy series (< 16 weeks)	Primary adult series (> 16 weeks)	Booster
Distemper	Respiratory, gastrointestinal and neurologic	1 dose at 6-8 weeks, then every 3-4 weeks until 16 weeks old	2 doses, 3-4 weeks apart	1 dose 1 year after primary series, then every 3 years
Adenovirus-2	Protective for ICH (hepatitis) virus	Same as above	Same as above	Same as above
Parvovirus	Gastrointestinal, highly contagious	Same as above	Same as above	Same as above
Parainfluenza	Respiratory	Same as above	Same as above	Same as above
Rabies	Neurologic, fatal, transmissible to humans	1 dose at 16 weeks	1 dose	1 dose 1 year after first dose, then every 3 years

Lifestyle Dependent Vaccines

Disease	Description	Primary puppy series	Primary adult series	Booster
Leptospirosis	Liver and kidney, transmissible to humans	2 doses, 2-4 weeks apart, beginning at 12 weeks	2 doses, 2-4 weeks apart	Annual
Bordetella + Parainfluenza	Kennel cough complex	1 dose at 4 weeks or older	1 dose	6-12 months, dependent on risk
Canine Influenza (H3N8)	Respiratory	2 doses, 3-4 weeks apart, at 12 weeks or older	2 doses, 3-4 weeks apart	Annual
Crotalus (rattlesnake)	Tissue damage, coagulation disorders	2 doses, 3-4 weeks apart, at 12 weeks or older	2 doses, 3-4 weeks apart	Annual